education

Department of Education
FREE STATE PROVINCE

PREPARATORY EXAMINATION



GRADE 12

PHYSICAL SCIENCES P2 (CHEMISTRY)



SEPTEMBER 2025

MARKS: 150

TIME: 3 HOURS

This question paper consists of 14 pages and 4 data sheets.

Copyright reserved

INSTRUCTIONS AND INFORMATION

1. Write your name and other applicable information in the appropriate spaces on the ANSWER BOOK.
2. The question paper consists of NINE questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two subquestions, e.g. between QUESTION 3.1 and QUESTION 3.2.
6. You may use a non-programmable pocket calculator.
7. You may use appropriate mathematical instruments.
8. Show ALL formulae and substitutions in ALL calculations.
9. Round off your FINAL numerical answers to a minimum of TWO decimal places where necessary.
10. Give brief motivations, discussions, etc. where required.
11. You are advised to use the attached DATA SHEETS.
12. Write neatly and legibly.

Copyright reserved

QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Each question has only ONE correct answer. Choose the answer and write only the letter (A–D) next to the question numbers (1.1 to 1.10) in the ANSWER BOOK, for example, 1.11 E.

1.1 Which ONE of the following is an unsaturated hydrocarbon?

A C_4H_{10}

B $CH_3(CH)_2CH_3$

C $CH_3(CH_2)_2CH_3$

D $CH_3CH_2CH_2CH_3$ (2)

1.2 Consider the organic molecule below.



Its correct IUPAC name is …


A butan-2-ol.

B butanoic acid.

C methyl butanoate.

D ethyl butanoate. (2)

1.3 Consider the organic structure below.



The IUPAC name of its FUNCTIONAL ISOMER is …

A methyl ethanoate.

B propanoic acid.

C ethyl methanoate.

D ethanoic acid. (2)

1.4 The reaction of copper(II) carbonate with EXCESS dilute hydrochloric acid is used to investigate the rate of reaction. The balanced equation for this reaction is:

$$CuCO_3(s) + 2HC\ell(aq) \rightarrow CuC\ell_2(aq) + H_2O(\ell) + CO_2(g)$$

The rate of reaction can experimentally be measured by measuring the …

A hydrochloric acid used.

B amount of water formed per time.

C volume of CO_2 produced per time.

D temperature of the mixture. (2)

1.5 Which ONE of the following statements concerning equilibrium is NOT correct?

A Equilibrium is reached in a closed system.

B At equilibrium, the concentration of reactants and products is constant.

C At equilibrium, the concentration of reactants and products is not necessarily the same.

D At equilibrium, the concentration of reactants is equal to the concentration of products. (2)

1.6 Write down the conjugate base of the following acid:

$$HC\ell O_4$$

A $HC\ell O_3^-$

B $HC\ell O_4^-$

C $C\ell O_4^-$

D $H_2C\ell O_4$ (2)

Stanmorephysics.com

1.7 The role of an indicator during a titration is to …

A show the color of the base.

B standardise the solution.

C show the color of the acid.

D identify the endpoint. (2)

1.8 Which statement is CORRECT for a Zn-Cu galvanic cell that operates under standard conditions?

A The concentration of the Zn^{2+} ions in the zinc half-cell gradually decreases.

B The concentration of the Cu^{2+} ions in the copper half-cell gradually increases.

C The intensity of the colour of the electrolyte in the copper half-cell gradually decreases.

D Negative ions migrate from the zinc half-cell to the copper half-cell. (2)

1.9 What is the energy convertion in an electrolytic cell?

A Electrical to chemical

B Chemical to electrical

C Electrical to mechanical

D Chemical to mechanical (2)

1.10 During the purification of copper, the …

A impure copper is the positive electrode.

B impure copper is connected to the cathode.



C pure copper is the positive electrode.

D pure copper is connected to the anode. (2)

[20]

QUESTION 2 (Start on a new page.)

The letters **A** to **F** in the table below represent organic compounds.

A	$CH_3COCH(CH_3)_2$	**B**	3,3-dimethylhexane
C		**D**	
E	$C_2H_4O_2$	**F**	Propan-1-ol

Stanmorephysics.com

2.1 Write down the letter(s) representing:

2.1.1 An aldehyde (1)

2.1.2 Functional isomers (1)

2.1.3 A carboxylic acid (1)

2.2 Write down the general formula of the homologous series to which compound **B** belongs. (2)

2.3 For compound **C**, write down the:

2.3.1 Letter of the molecule that is its structural isomer (1)

2.3.2 Type of structural isomerism named in QUESTION 2.2.1 (1)

2.3.3 IUPAC name (3)

2.4 For compound **A**, write down the structural formula of its chain isomer. (3)

2.5 Compounds **E** and **F** react to form an organic molecule. For this reaction, write down the:

2.5.1 Type of reaction (1)

2.5.2 NAME or FORMULA of the catalyst used (1)

2.5.3 IUPAC name of the ORGANIC product formed (2)

[17]

QUESTION 3 (Start on a new page.)

The boiling points of four organic compounds are shown in the table below.

	COMPOUND	BOILING POINT (°C)	MOLAR MASS ($g \cdot mol^{-1}$)
A	CH_3CH_2COOH	141,1	74
B	$CH_3(CH_2)_3OH$	….	74
C	$CH_3CH(CH_3)CH_2OH$	108	74
D	$CH_3COCH_2CH_3$	79,6	72

3.1 Define the term *boiling point*. (2)

3.2 Compounds **C's** boiling point is lower than that of compound **B**. Fully explain. (3)

3.3 Consider compounds **A**, **B** and **D**.

3.3.1 Predict the value of the boiling point of compound **B** in °C. Choose from:

145	141	117,7

(1)

3.3.2 Fully explain the answer to QUESTION 3.3.1. (6)

3.4 The vapour pressures of three straight-chained alkanes **E**, **F** and **G** are shown below.

COMPOUND		VAPOUR PRESSURE at 25 °C (kPa)
E	C_2H_6	4 187
F	C_3H_8	1 050
G	C_4H_{10}	250

3.4.1 Write the IUPAC NAME of the compound with the highest BOILING POINT. (2)


3.4.2 Fully explain the trend in the vapour pressure from compound **E** to compound **G**. (3)

[17]

Stanmorephysics.com

QUESTION 4 (Start on a new page.)

The flow diagram below shows a number of organic reactions. Reaction **I** is CATALYTIC CRACKING.



4.1 Define the term *cracking*. (2)

4.2 Consider reaction **II**.

4.2.1 What type of reaction is it? Choose from ENDOTHERMIC or EXOTHERMIC. (1)

4.2.2 Using MOLECULAR FORMULAE, write down the balanced chemical equation. (3)

4.2.3 Write down the NAME of compound **S**. (1)

4.3 In reaction **III**, compound **R** reacts with HBr to form compound **U**, which is a major product with molecular formula C_3H_7Br.

For reaction **III**, write down:

4.3.1 The type of reaction (1)

4.3.2 ONE reaction condition (1)

4.3.3 The IUPAC name of compound **U** (2)

4.4 For reaction **IV**, write down the:

4.4.1 Structural formula of the alcohol formed (2)

4.4.2 Type of reaction (1)

4.5 For reaction **V**, write down the:

4.5.1 Type of elimination reaction (1)

4.5.2 IUPAC name of compound **R** (2)
[17]

Stanmorephysics.com


QUESTION 5 (Start on a new page.)

A factor influencing the rate of a chemical reaction is investigated by conducting two experiments, **1** and **2,** in which the following reaction occurs:

$$Mg(s) + 2HC\ell(aq) \rightarrow MgC\ell_2(aq) + H_2(g)$$

In both experiments, the same mass of magnesium(Mg) is added to an EXCESS hydrochloric acid(HCℓ) solution with the same concentration in an open flask at STP.

The graph below shows the changes in the amount of gas collected during the reaction in experiments **1** and **2**.




5.1 Write down the name of the apparatus used for collecting the gas. (1)

5.2 Define the term *rate of reaction*. (2)

5.3 Calculate the average rate (in $dm^3 \cdot s^{-1}$) at which hydrogen is produced in experiment **1**. (3)

5.4 If 2 g of magnesium was used for this investigation, calculate the percentage purity of the magnesium. (7)

5.5 The Maxwell-Boltzmann distribution curves for the reaction in experiment **1** and **2** are given below.



5.5.1 Which factor affecting the reaction rate is being investigated? (1)

5.5.2 Which curve (**A** or **B**) represents the reaction in experiment **2**? (1)

5.5.3 Explain the answer to QUESTION 5.5.2 by referring to the collision theory. (2)
[17]

Stanmorephysics.com

QUESTION 6 (Start on a new page.)

Carbon, C, reacts with an unknown amount of carbon dioxide, CO_2, in a sealed 2 dm^3 container, at a temperature **T**. The equilibrium constant, K_c, at this temperature is 0,2. The balanced equation for the reaction is:

$$CO_2(g) + C(s) \rightleftharpoons 2CO(g)$$

At equilibrium, the concentration of carbon monoxide, CO, is found to be 0,04 $mol \cdot dm^{-3}$.

6.1 Can the above situation be referred to as a closed system?
Write down YES or NO. (1)

6.2 Explain the answer to QUESTION 6.1. (2)

6.3 Calculate the initial moles of CO_2 used. (8)

6.4 Is the yield of carbon monoxide HIGH or LOW? (1)

6.5 Explain the answer to QUESTION 6.4. (2)

6.6 When the temperature is decreased, the yield of CO decreases.

6.6.1 Is the forward reaction EXOTHERMIC or ENDOTHERMIC? (1)

6.6.2 Use Le Chatelier's principle to explain the answer to QUESTION 6.6.1. (2)

6.7 Draw a graph of potential energy ($kJ \cdot mol^{-1}$) versus the reaction coordinate for the reverse reaction.

Indicate the following on the graph:

- Activation energy
- Heat of reaction (ΔH)
- Energy of reactants and products

Write down the actual formulae of reactants and products (CO_2, C and CO) on the graph as labels of these energies. (4)
[21]

QUESTION 7 (Start on a new page.)

7.1 A standard solution of NaOH with a pH of 13,4 is used to neutralise 3,125 cm^3 of a 0,4 $mol \cdot dm^{-3}$ solution of H_2SO_4. The balanced chemical equation of the reaction is:

$$H_2SO_4(aq) + 2NaOH(aq) \rightarrow Na_2SO_4(aq) + 2H_2O(\ell)$$

7.1.1 Define the term *standard solution*. (2)

7.1.2 Is NaOH a strong or weak base? (1)

7.1.3 Explain the answer to QUESTION 7.1.2. (2)

7.1.4 Calculate the concentration of the NaOH solution. (5)

7.1.5 Calculate the volume of the NaOH needed to neutralise the acid. (5)

7.2 A laboratory technician neutralises an accidental spill by pouring 5 g of baking soda ($NaHCO_3$) on a 50 cm^3 hydrochloric acid solution of concentration 1,6 $mol \cdot dm^{-3}$.

Stanmorephysics.com

$$HC\ell(aq) + NaHCO_3(s) \rightarrow NaC\ell(aq) + CO_2(g) + H_2O(\ell)$$

Calculate the pH of the final solution. Assume the volume remains constant. (7)

[22]

QUESTION 8 (Start on a new page.)

A learner has the following reagents to construct an operational GALVANIC CELL under standard conditions: Lead, Pb, lead nitrate, $Pb(NO_3)_2$, silver, Ag, silver nitrate, $AgNO_3$, and iron(III) chloride, $FeC\ell_3$

8.1 Define the term *electrolyte*. (2)

8.2 Using ONLY the reagents given above, write down the cell notation of the operational galvanic cell. (3)


8.3 Write down the NAME or FORMULA of a substance that can be used in the salt bridge. (2)

8.4 Calculate the EMF of the above cell. (4)
[11]

Stanmorephysics.com

QUESTION 9 (Start on a new page.)

The simplified diagram below represents the electrolysis of a copper(II) chloride solution.



Stanmorephysics.com

9.1 Write down ONE safety measure which must be taken during the above process. (1)

9.2 Explain why the safety measure in QUESTION 9.1 is necessary. (2)

9.3 In what direction are the electrons moving? Choose from **P to Q** or **Q to P**. (1)

9.4 Write down the NAME of the substance that forms at electrode **P**. (1)

9.5 Write down the balanced net ionic equation for the reaction. (3)
[8]

TOTAL: 150

Copyright reserved

DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)

GEGEWENS VIR FISIESE WETENSKAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)

TABLE 1: PHYSICAL CONSTANTS / *TABEL 1: FISIESE KONSTANTES*

NAME / *NAAM*	SYMBOL / *SIMBOOL*	VALUE / *WAARDE*
Standard pressure *Standaarddruk*	p^θ	$1{,}013 \times 10^5$ Pa
Molar gas volume at STP *Molêre gasvolume by STD*	V_m	22,4 $dm^3 \cdot mol^{-1}$
Standard temperature *Standaardtemperatuur*	T^θ	273 K
Charge on electron *Lading op elektron*	e	$-1{,}6 \times 10^{-19}$ C
Avogadro's constant *Avogadro-konstante*	N_A	$6{,}02 \times 10^{23}$ mol^{-1}

TABLE 2: FORMULAE / *TABEL 2: FORMULES*

Stanmorephysics.com

$n=\frac{m}{M}$	$n=\frac{N}{N_A}$
$c=\frac{n}{V}$ *OR/OF* $c=\frac{m}{MV}$	$n=\frac{V}{V_m}$
$\frac{c_a V_a}{c_b V_b}=\frac{n_a}{n_b}$	$pH=-\log[H_3O^+]$
$K_w=[H_3O^+][OH^-]=1\times10^{-14}$ *at/by 298 K*	
$E^\theta_{cell}=E^\theta_{cathode}-E^\theta_{anode}$ *OR* $E^\theta_{cell}=E^\theta_{reduction}-E^\theta_{oxidation}$ *OR* $E^\theta_{cell}=E^\theta_{oxidising\ agent}-E^\theta_{reducing\ agent}$	$E^\theta_{sel}=E^\theta_{katode}-E^\theta_{anode}$ *OF* $E^\theta_{sel}=E^\theta_{reduksie}-E^\theta_{oksidasie}$ *OR* $E^\theta_{sel}=E^\theta_{oksideermiddel}-E^\theta_{reduseermiddel}$

Stanmorephysics.com

TABLE 3: THE PERIODIC OF ELEMENTS

1 (I)	2 (II)	3	4	5	6	7	8	9	10	11	12	13 (III)	14 (IV)	15 (V)	16 (VI)	17 (VII)	18 (VIII)
2,1 H 1																	2 He 4
1,0 Li 7	1,5 Be 9											2,0 B 11	2,5 C 12	3,0 N 14	3,5 O 16	4,0 F 19	10 Ne 20
0,9 Na 23	1,2 Mg 24											1,5 Aℓ 27	1,8 Si 28	2,1 P 31	2,5 S 32	3,0 Cℓ 35,5	18 Ar 40
0,8 K 39	1,0 Ca 40	1,3 Sc 45	1,5 Ti 48	1,6 V 51	1,6 Cr 52	1,5 Mn 55	1,8 Fe 56	1,8 Co 59	1,8 Ni 59	1,9 Cu 63,5	1,6 Zn 65	1,6 Ga 70	1,8 Ge 73	2,0 As 75	2,4 Se 79	2,8 Br 80	36 Kr 84
0,8 Rb 86	1,0 Sr 88	1,2 Y 89	1,4 Zr 91	Nb 92	1,8 Mo 96	1,9 Tc	2,2 Ru 101	2,2 Rh 103	2,2 Pd 106	1,9 Ag 108	1,7 Cd 112	1,7 In 115	1,8 Sn 119	1,9 Sb 122	2,1 Te 128	2,5 I 127	54 Xe 131
0,7 Cs 133	0,9 Ba 137	La 139	1,6 Hf 179	Ta 181	W 184	Re 186	Os 190	Ir 192	Pt 195	Au 197	Hg 201	1,8 Tℓ 204	1,8 Pb 207	1,9 Bi 209	2,0 Po	2,5 At	86 Rn
0,7 Fr	0,9 Ra 226	Ac															

58 Ce 140	59 Pr 141	60 Nd 144	61 Pm	62 Sm 150	63 Eu 152	64 Gd 157	65 Tb 159	66 Dy 163	67 Ho 165	68 Er 167	69 Tm 169	70 Yb 173	71 Lu 175
90 Th 232	91 Pa	92 U 238	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr

KEY/*SLEUTEL*

Atomic number
Atoomgetal

Electronegativity
Elektronegatiwiteit

Symbol
Simbool

Approximate relative atomic mass
Benaderde relatiewe atoommassa

Copyright reserved

TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE

Half-reactions / *Halfreaksies*	E^θ (V)
$F_2(g) + 2e^- \rightleftharpoons 2F^-$	+ 2,87
$Co^{3+} + e^- \rightleftharpoons Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^- \rightleftharpoons 2H_2O$	+1,77
$MnO_4^- + 8H^+ + 5e^- \rightleftharpoons Mn^{2+} + 4H_2O$	+ 1,51
$C\ell_2(g) + 2e^- \rightleftharpoons 2C\ell^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^- \rightleftharpoons 2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^- \rightleftharpoons 2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^- \rightleftharpoons Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^- \rightleftharpoons Pt$	+ 1,20
$Br_2(\ell) + 2e^- \rightleftharpoons 2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^- \rightleftharpoons NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^- \rightleftharpoons Hg(\ell)$	+ 0,85
$Ag^+ + e^- \rightleftharpoons Ag$	+ 0,80
$NO_3^- + 2H^+ + e^- \rightleftharpoons NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^- \rightleftharpoons Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^- \rightleftharpoons H_2O_2$	+ 0,68
$I_2 + 2e^- \rightleftharpoons 2I^-$	+ 0,54
$Cu^+ + e^- \rightleftharpoons Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^- \rightleftharpoons S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^- \rightleftharpoons 4OH^-$	+ 0,40
$Cu^{2+} + 2e^- \rightleftharpoons Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^- \rightleftharpoons SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^- \rightleftharpoons Cu^+$	+ 0,16
$Sn^{4+} + 2e^- \rightleftharpoons Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^- \rightleftharpoons H_2S(g)$	+ 0,14
$2H^+ + 2e^- \rightleftharpoons H_2(g)$	**0,00**
$Fe^{3+} + 3e^- \rightleftharpoons Fe$	– 0,06
$Pb^{2+} + 2e^- \rightleftharpoons Pb$	– 0,13
$Sn^{2+} + 2e^- \rightleftharpoons Sn$	– 0,14
$Ni^{2+} + 2e^- \rightleftharpoons Ni$	– 0,27
$Co^{2+} + 2e^- \rightleftharpoons Co$	– 0,28
$Cd^{2+} + 2e^- \rightleftharpoons Cd$	– 0,40
$Cr^{3+} + e^- \rightleftharpoons Cr^{2+}$	– 0,41
$Fe^{2+} + 2e^- \rightleftharpoons Fe$	– 0,44
$Cr^{3+} + 3e^- \rightleftharpoons Cr$	– 0,74
$Zn^{2+} + 2e^- \rightleftharpoons Zn$	– 0,76
$2H_2O + 2e^- \rightleftharpoons H_2(g) + 2OH^-$	– 0,83
$Cr^{2+} + 2e^- \rightleftharpoons Cr$	– 0,91
$Mn^{2+} + 2e^- \rightleftharpoons Mn$	– 1,18
$A\ell^{3+} + 3e^- \rightleftharpoons A\ell$	– 1,66
$Mg^{2+} + 2e^- \rightleftharpoons Mg$	– 2,36
$Na^+ + e^- \rightleftharpoons Na$	– 2,71
$Ca^{2+} + 2e^- \rightleftharpoons Ca$	– 2,87
$Sr^{2+} + 2e^- \rightleftharpoons Sr$	– 2,89
$Ba^{2+} + 2e^- \rightleftharpoons Ba$	– 2,90
$Cs^+ + e^- \rightleftharpoons Cs$	- 2,92
$K^+ + e^- \rightleftharpoons K$	– 2,93
$Li^+ + e^- \rightleftharpoons Li$	– 3,05

Increasing oxidising ability / *Toenemende oksiderende vermoë* ↑

Increasing reducing ability / *Toenemende reduserende vermoë* ↓

Copyright reserved

TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE

Half-reactions / *Halfreaksies*			E^{θ} (V)
$Li^+ + e^-$	$\rightleftharpoons$	Li	– 3,05
$K^+ + e^-$	$\rightleftharpoons$	K	– 2,93
$Cs^+ + e^-$	$\rightleftharpoons$	Cs	– 2,92
$Ba^{2+} + 2e^-$	$\rightleftharpoons$	Ba	– 2,90
$Sr^{2+} + 2e^-$	$\rightleftharpoons$	Sr	– 2,89
$Ca^{2+} + 2e^-$	$\rightleftharpoons$	Ca	– 2,87
$Na^+ + e^-$	$\rightleftharpoons$	Na	– 2,71
$Mg^{2+} + 2e^-$	$\rightleftharpoons$	Mg	– 2,36
$A\ell^{3+} + 3e^-$	$\rightleftharpoons$	$A\ell$	– 1,66
$Mn^{2+} + 2e^-$	$\rightleftharpoons$	Mn	– 1,18
$Cr^{2+} + 2e^-$	$\rightleftharpoons$	Cr	– 0,91
$2H_2O + 2e^-$	$\rightleftharpoons$	$H_2(g) + 2OH^-$	– 0,83
$Zn^{2+} + 2e^-$	$\rightleftharpoons$	Zn	– 0,76
$Cr^{3+} + 3e^-$	$\rightleftharpoons$	Cr	– 0,74
$Fe^{2+} + 2e^-$	$\rightleftharpoons$	Fe	– 0,44
$Cr^{3+} + e^-$	$\rightleftharpoons$	Cr^{2+}	– 0,41
$Cd^{2+} + 2e^-$	$\rightleftharpoons$	Cd	– 0,40
$Co^{2+} + 2e^-$	$\rightleftharpoons$	Co	– 0,28
$Ni^{2+} + 2e^-$	$\rightleftharpoons$	Ni	– 0,27
$Sn^{2+} + 2e^-$	$\rightleftharpoons$	Sn	– 0,14
$Pb^{2+} + 2e^-$	$\rightleftharpoons$	Pb	– 0,13
$Fe^{3+} + 3e^-$	$\rightleftharpoons$	Fe	– 0,06
$2H^+ + 2e^-$	**$\rightleftharpoons$**	**$H_2(g)$**	**0,00**
$S + 2H^+ + 2e^-$	$\rightleftharpoons$	$H_2S(g)$	+ 0,14
$Sn^{4+} + 2e^-$	$\rightleftharpoons$	Sn^{2+}	+ 0,15
$Cu^{2+} + e^-$	$\rightleftharpoons$	Cu^+	+ 0,16
$SO_4^{2-} + 4H^+ + 2e^-$	$\rightleftharpoons$	$SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + 2e^-$	$\rightleftharpoons$	Cu	+ 0,34
$2H_2O + O_2 + 4e^-$	$\rightleftharpoons$	$4OH^-$	+ 0,40
$SO_2 + 4H^+ + 4e^-$	$\rightleftharpoons$	$S + 2H_2O$	+ 0,45
$Cu^+ + e^-$	$\rightleftharpoons$	Cu	+ 0,52
$I_2 + 2e^-$	$\rightleftharpoons$	$2I^-$	+ 0,54
$O_2(g) + 2H^+ + 2e^-$	$\rightleftharpoons$	H_2O_2	+ 0,68
$Fe^{3+} + e^-$	$\rightleftharpoons$	Fe^{2+}	+ 0,77
$NO_3^- + 2H^+ + e^-$	$\rightleftharpoons$	$NO_2(g) + H_2O$	+ 0,80
$Ag^+ + e^-$	$\rightleftharpoons$	Ag	+ 0,80
$Hg^{2+} + 2e^-$	$\rightleftharpoons$	$Hg(\ell)$	+ 0,85
$NO_3^- + 4H^+ + 3e^-$	$\rightleftharpoons$	$NO(g) + 2H_2O$	+ 0,96
$Br_2(\ell) + 2e^-$	$\rightleftharpoons$	$2Br^-$	+ 1,07
$Pt^{2+} + 2e^-$	$\rightleftharpoons$	Pt	+ 1,20
$MnO_2 + 4H^+ + 2e^-$	$\rightleftharpoons$	$Mn^{2+} + 2H_2O$	+ 1,23
$O_2(g) + 4H^+ + 4e^-$	$\rightleftharpoons$	$2H_2O$	+ 1,23
$Cr_2O_7^{2-} + 14H^+ + 6e^-$	$\rightleftharpoons$	$2Cr^{3+} + 7H_2O$	+ 1,33
$C\ell_2(g) + 2e^-$	$\rightleftharpoons$	$2C\ell^-$	+ 1,36
$MnO_4^- + 8H^+ + 5e^-$	$\rightleftharpoons$	$Mn^{2+} + 4H_2O$	+ 1,51
$H_2O_2 + 2H^+ + 2e^-$	$\rightleftharpoons$	$2H_2O$	+1,77
$Co^{3+} + e^-$	$\rightleftharpoons$	Co^{2+}	+ 1,81
$F_2(g) + 2e^-$	$\rightleftharpoons$	$2F^-$	+ 2,87

Increasing oxidising ability / *Toenemende oksiderende vermoë* ↓

Increasing reducing ability / *Toenemende reduserende vermoë* ↑

Copyright reserved